

# World Journal of Pharmaceutical Sciences



[Register](#) [Login](#)



**ISSN (Print): 2321-3310**

**ISSN (Online): 2321-3086**

**US CODEN : WJPSBT**

**\* Impact Factor: 1.877 (2013)**

- 
- [Home](#)
- [Editorial Desk](#)
- [Current Issue](#)
- [Authors Instructions](#)
- [Manuscript Submission](#)
- [Downloads](#)
- [Contact Us](#)



**What's New:**



## World Journal of Pharmaceutical Sciences



Register (login.php)

or



Login (login.php)

ISSN (Print): 2321-3310

ISSN (Online): 2321-3086

US CODEN: WJPSBT

Impact Factor: 0.453 (2015)

MENU



Issue: [World J Pharm Sci 2015; 3(7) : 1283-1470]

**Editorial: Targeted therapy as an effective strategy in the superb modern science and technology era**  
(view\_issue.php?title=Targeted-therapy-as-an-effective-strategy-in-the-superb-modern-science-and-technology-era)

**Dr. Behzad Foroutan**

Pages: 1283-1285

Abstract

Full Text (admin/uploads/sn3UTX.pdf)

Download PDF (download.php?id=NDg0)

XML </> (issue.xml?title=Targeted-therapy-as-an-effective-strategy-in-the-superb-modern-science-and-technology-era)

Views (843) ()

Downloads (606) ()

**Editorial: Emerging role of herbal drugs in the management of diabetes mellitus**  
(view\_issue.php?title=Emerging-role-of-herbal-drugs-in-the-management-of-diabetes-mellitus)

**Harsimran Singh, Jagdeep Kaur, Nitish Kumar, MU Khan**

Pages: 1286-1289

Abstract

Full Text (admin/uploads/kz9ldG.pdf)

Download PDF (download.php?id=NDg1)

XML </> (issue.xml?title=Emerging-role-of-herbal-drugs-in-the-management-of-diabetes-mellitus)

Views (817) ()

Downloads (725) ()

**Original Article: Simultaneous determination of tapentadol HCl and paracetamol by ratio-spectra derivative spectrophotometry**  
(view\_issue.php?title=Simultaneous-determination-of-tapentadol-HCl-and-paracetamol-by-ratio-spectra-derivative-spectrophotometry)

**Hamed M. El-Fatraty, Mokhtar M. Mabrouk, Sherin F. Hammad, and Samah F. El-Malla**

Pages: 1290-1297

Abstract

Full Text (admin/uploads/PT930J.pdf)

Download PDF (download.php?id=NDg2)

XML </> (issue.xml?title=Simultaneous-determination-of-tapentadol-HCl-and-paracetamol-by-ratio-spectra-derivative-spectrophotometry)

Views (733) ()

Downloads (599) ()

**Original Article: Evaluation of the Knowledge and Practice of Pharmaceutical Care among Hospital Pharmacists in Secondary and Tertiary Hospitals in Lagos State, Nigeria**  
(view\_issue.php?title=Evaluation-of-the-Knowledge-and-Practice-of-Pharmaceutical-Care-among-Hospital-Pharmacists-in-Secondary-and-Tertiary-Hospitals-in-Lagos-State-Nigeria)

**Eniojukan J Funsho and Onedo O Titilayo**

Pages: 1298-1306

Abstract

Full Text (admin/uploads/v213P4.pdf)

Download PDF (download.php?id=NDg3)

XML </> (issue.xml?title=Evaluation-of-the-Knowledge-and-Practice-of-Pharmaceutical-Care-among-Hospital-Pharmacists-in-Secondary-and-Tertiary-Hospitals-in-Lagos-State-Nigeria)

Views (751) ()

Downloads (706) ()

**Original Article: Microbial transformation of some simple  $\beta$ -carboline alkaloids**  
(view\_issue.php?title=Microbial-transformation-of-some-simple-946-carboline-alkaloids)

Duaa E. Mohamed, Abdel  Rahim S. Ibrahim, Souzan M. Ibrahim, Amal M. Kabbash

Pages: 1307-1318

Abstract 

Full Text  (admin/uploads/x4Aa8X.pdf)

Download PDF  (download.php?id=NDg4)

XML </> (issue.xml?title=Microbial-transformation-of-some-simple-946-carboline-alkaloids)

Views (671) ()

Downloads (540) ()

**Original Article: Knowledge, practice and complications associated with diabetic foot**  
(view\_issue.php?title=Knowledge-practice-and-complications-associated-with-diabetic-foot)

Fakhsheena Anjum, Sana Ghayas, Noor Jahan, Hina Yasin, M. Ali Masood

Pages: 1319-1323

Abstract 

Full Text  (admin/uploads/g982FK.pdf)

Download PDF  (download.php?id=NDg5)

XML </> (issue.xml?title=Knowledge-practice-and-complications-associated-with-diabetic-foot)

Views (739) ()


Downloads (503) ()

**Original Article: Synthesis, molecular docking and antitumor activity of N,N'-Carbonylbis(N-Ethylbenzamide)**  
(view\_issue.php?title=Synthesis-molecular-docking-and-antitumor-activity-of-NN-CarbonylbisN-Ethylbenzamide-)

Nuzul Wahyuning Diyah, **Bambang Tri Purwanto**, Siswandono

Pages: 1324-1329

Abstract 

Full Text  (admin/uploads/yaPWdK.pdf)

Download PDF  (download.php?id=NDkw)

XML </> (issue.xml?title=Synthesis-molecular-docking-and-antitumor-activity-of-NN-CarbonylbisN-Ethylbenzamide-)

Views (902) ()


Downloads (675) ()

**Original Article: Gradual Decline in the Potency of Antipseudomonas Drugs: Evaluated by Retrospective Comparative studies of Susceptibility and Resistance Pattern of Antimicrobials against Pseudomonas Aeruginosa Isolates Obtained from Intensive Care Unit**  
(view\_issue.php?title=Gradual-Decline-in-the-Potency-of-Antipseudomonas-Drugs-Evaluated-by-Retrospective-Comparative-studies-of-Susceptibility-and-Resistance-Pattern-of-Antimicrobials-against-Pseudomonas-Aeruginosa-Isolates-Obtained-from-Intensive-Care-Unit)

Somia Gul and Maria Ayub

Pages: 1330-1339

Abstract 

Full Text  (admin/uploads/Gyas3t.pdf)

Download PDF  (download.php?id=NDkx)

XML </> (issue.xml?title=Gradual-Decline-in-the-Potency-of-Antipseudomonas-Drugs-Evaluated-by-Retrospective-Comparative-studies-of-Susceptibility-and-Resistance-Pattern-of-Antimicrobials-again

Views (697) ()

Downloads (793) ()

**Original Article: Immunogenetic and Bacteriological Study of Acute Otitis Media and Tonsillitis among Children Patients**  
(view\_issue.php?title=Immunogenetic-and-Bacteriological-Study-of-Acute-Otitis-Media-and-Tonsillitis-among-Children-Patients-)

Abbas Zeki Mehdi, Hammadi A. Al-Hilali, Mohammed Sh. Jebur

Pages: 1340-1346

Abstract 

Full Text  (admin/uploads/YbN0GO.pdf)

Download PDF  (download.php?id=NDky)

XML </> (issue.xml?title=Immunogenetic-and-Bacteriological-Study-of-Acute-Otitis-Media-and-Tonsillitis-among-Children-Patients-)

Views (758) ()




Downloads (723) ()

**Original Article: Carboxymethyl cellulose based multifunctional targeted drug delivery platform for pancreatic cancer: Nanotheranostic potential and biocompatibility analysis**

(view\_issue.php?title=Carboxymethyl-cellulose-based-multifunctional-targeted-drug-delivery-platform-for-pancreatic-cancer-Nanotheranostic-potential-and-biocompatibility-analysis)

**Preeti Nigam Joshi, Shishanka Wangnoo, Michelle Louis**

Pages: 1347-1359




Abstract 
Full Text  (admin/uploads/xkTC2E.pdf)
Download PDF  (download.php?id=NDkz)
XML </> (issue.xml?title=Carboxymethyl-cellulose-based-multifunctional-targeted-drug-delivery-platform-for-pancreatic-cancer-Nanotheranostic-potential-and-biocompatibility-analysis)
Views (789) ()
Downloads (806) ()

**Original Article: Preliminary Phytochemical Screening, Anti-oxidant activity, Multi elemental analysis by ICP- Spectroscopy and Antimicrobial activity of bulbs extracts of Crinum woodrowii Baker from Bhimashankar, Maharashtra, India**

(view\_issue.php?title=Preliminary-Phytochemical-Screening-Anti-oxidant-activity-Multi-elemental-analysis-by-ICP-Spectroscopy-and-Antimicrobial-activity-of-bulbs-extracts-of-Crinum-woodrowii-Baker-from-Bhimashankar-Maharashtra-India)

**Ramjan Mulani and Sanjay Jagtap**

Pages: 1360-1369




Abstract 
Full Text  (admin/uploads/z8LbZf.pdf)
Download PDF  (download.php?id=NDk0)
XML </> (issue.xml?title=Preliminary-Phytochemical-Screening-Anti-oxidant-activity-Multi-elemental-analysis-by-ICP-Spectroscopy-and-Antimicrobial-activity-of-bulbs-extracts-of-Crinum-woodrowii-Baker-from-Bhimashankar-Maharashtra-India)
Views (676) ()
Downloads (538) ()

**Original Article: Quantitative and qualitative bacterial analysis of underground water of Vijayapur**

(view\_issue.php?title=Quantitative-and-qualitative-bacterial-analysis-of-underground-water-of-Vijayapur)

**Satihal S. S, Magadam S.R and Yadawe M. S**

Pages: 1370-1374




Abstract 
Full Text  (admin/uploads/O2JXCW.pdf)
Download PDF  (download.php?id=NDk1)
XML </> (issue.xml?title=Quantitative-and-qualitative-bacterial-analysis-of-underground-water-of-Vijayapur)
Views (744) ()
Downloads (475) ()

**Short Communication: A study on diabetic type 2 prevalence among urban population compared with rural population of Warangal and Adjoining Districts of Telangana**

(view\_issue.php?title=A-study-on-diabetic-type-2-prevalence-among-urban-population-compared-with-rural-population-of-Warangal-and-Adjoining-Districts-of-Telangana)

**Sireesha T, Asha S, Malathi J, Raghu Ram Reddy A, Mounika A, Mamatha P**

Pages: 1375-1380




Abstract 
Full Text  (admin/uploads/AUZy4B.pdf)
Download PDF  (download.php?id=NDk2)
XML </> (issue.xml?title=A-study-on-diabetic-type-2-prevalence-among-urban-population-compared-with-rural-population-of-Warangal-and-Adjoining-Districts-of-Telangana)
Views (711) ()
Downloads (465) ()

**Original Article: Assay of antibacterial agents against drug resistant and drug sensitive bacteria and identification of biologically active principles from Ceriops tagal stem extracts**

(view\_issue.php?title=Assay-of-antibacterial-agents-against-drug-resistant-and-drug-sensitive-bacteria-and-identification-of-biologically-active-principles-from-Ceriops-tagal-stem-extracts)

**S. Sunita, P. Satya Veni, and A. Srinivasulu**





Pages: 1381-1386

Abstract 
Full Text  (admin/uploads/14jEZR.pdf)
Download PDF  (download.php?id=NDk3)
XML </> (issue.xml?title=Assay-of-antibacterial-agents-against-drug-resistant-and-drug-sensitive-bacteria-and-identification-of-biologically-active-principles-from-Ceriops-tagal-stem-extracts)
Views (737) ()
Downloads (479) ()

**Original Article: Confidence, ease of use, satisfaction and preference of spacers in patients with asthma and COPD: Results from three open-label, prospective studies**  
(view\_issue.php?title=Confidence-ease-of-use-satisfaction-and-preference-of-spacers-in-patients-with-asthma-and-COPD-Results-from-three-open-label-prospective-studies)

**Sujeet Rajan, Sundeep Salvi, Jaideep Gogtay**





Pages: 1387-1396

Abstract 
Full Text  (admin/uploads/XL0n7Q.pdf)
Download PDF  (download.php?id=NDk4)
XML  (issue.xml?title=Confidence-ease-of-use-satisfaction-and-preference-of-spacers-in-patients-with-asthma-and-COPD-Results-from-three-open-label-prospective-studies)
Views (723) ()
Downloads (608) ()

**Original Article: Perception of patients towards the knowledge of community pharmacist as a health care provider: A pilot study**  
(view\_issue.php?title=Perception-of-patients-towards-the-knowledge-of-community-pharmacist-as-a-health-care-provider-A-pilot-study)

**Singh Dipti and Khale Anubha**





Pages: 1397-1402

Abstract 
Full Text  (admin/uploads/bqrcdG.pdf)
Download PDF  (download.php?id=NDk5)
XML  (issue.xml?title=Perception-of-patients-towards-the-knowledge-of-community-pharmacist-as-a-health-care-provider-A-pilot-study)
Views (776) ()
Downloads (512) ()

**Original Article: Pharmacognostic evaluation of medicinally important mangroves *Avicennia marina* and *Sonneratia apetala***  
(view\_issue.php?title=Pharmacognostic-evaluation-of-medicinally-important-mangroves-Avicennia-marina-and-Sonneratia-apetala-)

**Sunita Shailajan and Mayuresh Joshi**





Pages: 1403-1412

Abstract 
Full Text  (admin/uploads/XDLkP5.pdf)
Download PDF  (download.php?id=NTAw)
XML  (issue.xml?title=Pharmacognostic-evaluation-of-medicinally-important-mangroves-Avicennia-marina-and-Sonneratia-apetala-)
Views (682) ()
Downloads (772) ()

**Original Article: Isolation and characterization of a bioactive lectin from *Zizyphus oenoplia***  
(view\_issue.php?title=Isolation-and-characterization-of-a-bioactive-lectin-from-Zizyphus-oenoplia)

**Ashwin B. Butle and M. B. Patil**





Pages: 1413-1420

Abstract 
Full Text  (admin/uploads/UbNKG6.pdf)
Download PDF  (download.php?id=NTAx)
XML  (issue.xml?title=Isolation-and-characterization-of-a-bioactive-lectin-from-Zizyphus-oenoplia)
Views (752) ()
Downloads (619) ()

**Original Article: Comparative qualitative analysis of callus extracts of in-vitro and in-vivo plants of *Jasminum angustifolium*, a wild and medicinal plant**  
(view\_issue.php?title=Comparative-qualitative-analysis-of-callus-extracts-of-in-vitro-and-in-vivo-plants-of-Jasminum-angustifolium-a-wild-and-medicinal-plant)

**Painthamizharasi Lakshmanan and Jeya Jothi Gabriel**

Pages: 1421-1425



Abstract 
Full Text  (admin/uploads/LVRUPB.pdf)
Download PDF  (download.php?id=NTAy)
XML  (issue.xml?title=Comparative-qualitative-analysis-of-callus-extracts-of-in-vitro-and-in-vivo-plants-of-Jasminum-angustifolium-a-wild-and-medicinal-plant)
Views (712) ()
Downloads (502) ()

**Original Article: Phytochemical screening and estimation of total phenolics and total flavonoid content of *Cocos nucifera* endocarp**  
(view\_issue.php?title=Phytochemical-screening-and-estimation-of-total-phenolics-and-total-flavonoid-content-of-Cocos-nucifera-endocarp)

**Nidhi Tyagi, Vikas Hooda and Sachin malkani**

Pages: 1426-1432




Abstract 
--

Full Text  (admin/uploads/0Qv7jH.pdf)
Download PDF  (download.php?id=NTAz)
XML </> (issue.xml?title=Phytochemical-screening-and-estimation-of-total-phenolics-and-total-flavonoid-content-of-Cocos-nucifera-endocarp)
Views (721) ()
Downloads (622) ()

**Original Article: Formulation, characterization and evaluation of bioadhesive buccal patch of venlafaxine**  
(view\_issue.php?title=Formulation-characterization-and-evaluation-of-bioadhesive-buccal-patch-of-venlafaxine)

Sumit Durgapal, Sayantan Mukhopadhyay, Laxmi Goswami




Pages: 1433-1445

Abstract 
Full Text  (admin/uploads/38Jj57.pdf)
Download PDF  (download.php?id=NTA0)
XML </> (issue.xml?title=Formulation-characterization-and-evaluation-of-bioadhesive-buccal-patch-of-venlafaxine)
Views (684) ()
Downloads (468) ()

**Original Article: Comparison of alkaline phosphatase assay using two different buffer systems: AMP and DEA**  
(view\_issue.php?title=Comparison-of-alkaline-phosphatase-assay-using-two-different-buffer-systems-AMP-and-DEA)

Rajeswari S, Gandhi M, Arun Kumar D, Dhananjayan R and Swaminathan S




Pages: 1446-1449

Abstract 
Full Text  (admin/uploads/jbGk3t.pdf)
Download PDF  (download.php?id=NTA1)
XML </> (issue.xml?title=Comparison-of-alkaline-phosphatase-assay-using-two-different-buffer-systems-AMP-and-DEA)
Views (1174) ()
Downloads (650) ()

**Original Article: Purification and characterization of  $\beta$ -galactosidase from leaves of Zizyphus oenoplia**  
(view\_issue.php?title=Purification-and-characterization-of- $\beta$ -galactosidase-from-leaves-of-Zizyphus-oenoplia)

Ashwin Butle and M. B. Patil




Pages: 1450-1455


Abstract 
Full Text  (admin/uploads/2vkjJz.pdf)
Download PDF  (download.php?id=NTA2)
XML </> (issue.xml?title=Purification-and-characterization-of- $\beta$ -galactosidase-from-leaves-of-Zizyphus-oenoplia)
Views (680) ()
Downloads (500) ()

**Review: Current prospective of magnetically triggered microsphere towards colonic disease: A review**  
(view\_issue.php?title=Current-prospective-of-magnetically-triggered-microsphere-towards-colonic-disease-A-review)

Surender Verma, Neha Khar, Lokesh Sadana




Pages: 1456-1464

Abstract 
Full Text  (admin/uploads/5mH6k3.pdf)
Download PDF  (download.php?id=NTA3)
XML </> (issue.xml?title=Current-prospective-of-magnetically-triggered-microsphere-towards-colonic-disease-A-review)
Views (794) ()
Downloads (600) ()

**Original Article: Enzymatic and non-enzymatic antioxidant properties of Indigofera longeracemosa  An in vitro study**  
(view\_issue.php?title=Enzymatic-and-non-enzymatic-antioxidant-properties-of-Indigofera-longeracemosa-An-in-vitro-study)

V. Suseela, A. Dharani and R. Suganya

Pages: 1465-1470

Abstract 
Full Text  (admin/uploads/rLfbPZ.pdf)
Download PDF  (download.php?id=NTA4)
XML </> (issue.xml?title=Enzymatic-and-non-enzymatic-antioxidant-properties-of-Indigofera-longeracemosa-An-in-vitro-study)
Views (699) ()
Downloads (831) ()



## Synthesis, molecular docking and antitumor activity of *N,N'*-Carbonylbis(*N*-Ethylbenzamide)

Nuzul Wahyuning Diyah, Bambang Tri Purwanto, Siswandono

Pharmaceutical Chemistry Departement, Faculty of Pharmacy Airlangga University, Surabaya - Indonesia

Received: 24-05-2015 / Revised: 19-06-2015 / Accepted: 27-06-2015

### ABSTRACT

We have designed and synthesized *N,N'*-carbonylbis(*N*-ethylbenzamide) to find new urea derivative with antitumor activity. The compound was synthesized from reaction of *N,N'*-diethylurea and benzoyl chloride in tetrahydrofuran. The designed molecule was docked into binding site of p38 $\alpha$  MAP Kinase (pdb. 3HEG) to predict its binding affinity. The antitumor activity was tested *in vitro* on human breast cancer cell line (MCF-7 and T47D) by MTT assay and compared with hydroxyurea as a reference compound. The synthesis yield 30% and confirmed as *N,N'*-carbonylbis(*N*-ethylbenzamide) by spectroscopic data. The compound showed higher *in vitro* antitumor activity than hydroxyurea either against MCF-7 or T47D and displayed better *in silico* binding property. It concluded that the synthesized compound is recommended to be developed further antitumor agents for human breast cancer.

**Keywords:** *N,N'*-carbonylbis(*N*-ethylbenzamide), synthesis, 3HEG, antitumor activity.



### INTRODUCTION

Cancer is one of the leading causes of morbidity and mortality worldwide, with approximately 14 million new cases and 8.2 million cancer-related deaths in 2012 [1]. The number of new cases is expected to rise by about 70% over the next 2 decades. More than 60% of world's total new annual cases occur in Africa, Asia and Central and South America. These regions account for 70% of the world's cancer deaths. Among women the 5 most common sites diagnosed were breast, colorectum, lung, cervix, and stomach cancer. Meanwhile, anticancer drugs that have long been used gradually becomes less effective and the cancer cells becomes resistant to anticancer drugs [2, 3, 4]. Therefore, there is an urgent need for development of new anticancer agents with divergent and unique structure which is different from the existing antitumor agents.

It has been well established that urea derivatives have got a significant place in modern medicinal chemistry. Urea derivatives have been reported in the literature as insecticide [5], anticonvulsant [6], antibacterial and antifungal agents [7], anticancer agent [8], hypoglycemic agent [9]. Furthermore, a large variety of urea derivatives were reported to

possess potent inhibiting effects on receptor tyrosine kinases (RTKs) [10], raf-1 kinases [11], p38 $\alpha$  MAP kinases [12]. The p38 mitogen-activated protein kinase (p38 MAPK) is a crucial component for the regulation of many cellular processes, including cell proliferation, differentiation, survival and migration [13]. Recent evidence has shown that p38 $\alpha$  may also play a causative role in cancer.

Hydroxyurea (HU) is a classical anticancer drugs which is still usefull to treat several types of cancer, but not including breast cancer [14, 15, 16]. Nevertheless, it has been reported that HU has antitumor activity on breast cancer cell line with IC<sub>50</sub> of 307  $\mu$ M [17]. Another urea derivative, sorafenib, is an inhibitor of several tyrosine kinase and Raf kinase [18]. Nambodiri et al. (2010) reported that sorafenib could interacted with p38 $\alpha$  MAPK [19]. *In silico* study indicated that N-CO-N in urea's structures are pharmacophores that capable to form hydrogen bonding with amino acid in binding site of the target enzymes [20].

A serie of ring-substituted benzoylurea derivatives had reported were more active than hydroxyurea according to Brine Shrimp Lethality Test. Quantitative Structure Activity Relationship and *in*

*silico* study revealed that binding affinity of benzoylurea derivatives to the enzyme were better than hydroxyurea due to steric (MR) and hydrophobic (log P) factors and the presence of aromatic ring in the structure of urea [21]. In subsequent study, benzoylurea has been modified into *N,N'*-dibenzoyl-*N,N'*-dimethylurea which exhibits antitumor activity against human breast cancer cell line, MCF-7 [22]. Methyl substitution on both the N urea and one additional aromatic ring on the other N increase the activity of benzoylurea. Antiproliferative activity test using MTT assay was reported by many researcher [23, 24]. El-Shawy *et al.* (2011) and Lu *et al.* (2013) reported that aromatic ring and urea group in benzylurea is the pharmacophore responsible to tumor growth inhibition activity of some benzylurea derivatives [25, 26].

In the present study, we attached two ethyl groups and two aromatic rings on both N atoms of urea to obtain *N,N'*-carbonylbis(*N*-ethylbenzamide). The synthesis procedure was based on Schotten-Baumann acylation method [27], with modification in solvent. Antitumor activity were tested using MTT assay against two human breast cancer cell line, T47D and MCF-7, and compared with two anticancer drugs, hydroxyurea which containing same pharmacophore and 5-fluorouracil which has been clinically used in the treatment of breast cancer. *In-silico* activity were performed by Molegro Virtual Docker 5.5 using 3HEG as target molecule.

## MATERIALS AND METHODS

**General:** All reagents and solvents were purchased from standard commercial suppliers. *N,N'*-carbonylbis(*N*-ethylbenzamide) (**Figure 1**) were synthesized from the reaction of starting material *N,N'*-diethylurea with benzoyl chlorides in the organic solvent, tetrahydrofuran (THF), and using triethylamine as catalyst. The structure of synthesized compound were identified by IR, <sup>1</sup>H-NMR, <sup>13</sup>C-NMR, and MS spectroscopy; whereas their purity were determined by melting point and TLC tests.

Melting point was measured with an Electrothermal melting point apparatus without correction. IR spectrum was recorded in KBr on Jasco FT-IR 5300 and major absorption was listed in cm<sup>-1</sup>. <sup>1</sup>H-NMR and <sup>13</sup>C-NMR spectrum was taken at JEOL ECS-400 spectrometer (400MHz), and chemical shifts were reported in ppm on the  $\delta$ -scale from internal standard Me<sub>4</sub>Si. Mass spectrum was measured with Agilent 6890N spectrometer using EI method. TLC was carried out on aluminium plates coated with silica gel F<sub>254</sub> (E

Merck) using UV lamp 254 nm to spot detection. *In silico* molecular docking was running out using Molegro Virtual Docker (MVD) ver 5.0 (CLC Bio). CS ChemBioDraw Ultra ver 11.0 (Cambridge Soft) was employed to built 3D structure of compounds. Antitumor activity was tested *in vitro* against MCF-7 and T47D cell line by MTT method and expressed in IC<sub>50</sub>, concentration of the compounds inducing a 50% inhibition of cell growth of treated cells compared to the growth of control cells. Hydroxyurea (HU) and 5-fluorouracil (5-FU) were used as the reference drugs. The absorbance at 595 nm was recorded using ELISA Microplate Reader (Benchmark). The IC<sub>50</sub> values were calculated by probit analysis using SPSS 20.

**Synthesis of *N,N'*-carbonylbis (*N*-ethylbenzamide):** *N,N'*-diethylurea 0,0125 mol dissolved in 20 ml THF was mixed with 4 ml triethylamine in a 200 ml conical flask. The solution of benzoyl chloride 0,0250 mol in 20 ml THF then added drop wise to the mixture at 0–5 °C ice bath while stirring. Then the temperature was raised to 70 °C and the mixture was refluxed for 3 hours. Reflux was continued at room temperature for next 20 hours and the mixture was concentrated by evaporating the solvent. The product was washed with cold water and saturated sodium bicarbonate respectively, and the solid residu was crystallized from ethanol–water (1:1).

***In silico* Molecular Docking:** The structure of p38 $\alpha$  MAP kinase (pdb. 3HEG at 2.2 Å resolution) was obtained from the Protein Data Bank ([www.rcsb.org](http://www.rcsb.org)) [28]. Compound structures were built with ChemBioDraw Ultra 11.0 and their geometry optimization were performed using the MMFF94 method in the program and saved as Sybyl Mol2 format. In the molecular docking, the test compounds were placed into binding site of 3HEG (cavity-1) by align method to the reference ligand (BAX). BAX is sorafenib molecule, an urea anticancer drug which capable to complex with the enzyme [19]. The binding affinity between ligand and enzyme (docking score) was predicted using MolDock Score and post analysed as Rerank Score (RS). The highest-scoring pose (lowest energy) should represent the best-found binding mode. Evaluation of the interaction was based on the their RS which is the sum of ligand-protein interaction energy and internal ligand energy, including hydrogen bonds between ligands and protein. The validation of docking was carried out by redocking the BAX into binding site (cavity-1) of 3HEG. A more negative binding affinity indicates stronger binding. The best docked structures have to follow these criteria : i) they must occupy the same cavity in the enzyme similar to BAX, (ii) they have the highest-scoring pose (lowest binding energy);



which can be observed visually by comparing the structure of docked molecule with crystal structure of BAX inside the binding site. The ligand-protein complexes of the top-scoring poses were used for further visual inspection.

**In vitro Antitumor activity:** In vitro antitumor activity was tested by MTT assay and expressed in  $IC_{50}$ , concentration of the compounds inducing a 50% inhibition of cell growth of treated cells compared to the growth of untreated cells [24]. Hydroxyurea and 5-Fluorouracil used as reference drugs. Cells were seeded at a density of  $8 \times 10^4$  cells/well into 96-well microplates, followed by treatment with tested compounds at concentration ranging between 100 – 1000  $\mu\text{g/mL}$  and incubated in 5%  $\text{CO}_2$  incubator at  $37^\circ\text{C}$ . After 24 hours, each wells were added by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) solution (100  $\mu\text{L}$  of 0.5  $\text{mg/mL}$  MTT in PBS). Then 100  $\mu\text{L}$  10% SDS in 0.1 N HCl was added after 3 hours and plates were incubated in the dark at  $37^\circ\text{C}$  overnight. Cell viability was determined from the absorbance which was observed at 595 nm on ELISA-reader and expressed in percentages with respect to untreated cells. The  $IC_{50}$  values was calculated with probit analysis. Each experiment was performed at least three times.

## RESULT AND DISCUSSION

### Synthesis product, *N,N'*-carbonylbis (*N*-ethylbenzamide):

*N,N'*-carbonylbis(*N*-ethylbenzamide) (Figure 1) were synthesized from *N,N'*-diethylurea in one steps. The compounds is white solids and water insoluble substances. Structure of the compound were identified by IR,  $^1\text{H-NMR}$ ,  $^{13}\text{C-NMR}$ , and MS spectroscopy as follows : Yield 30% as white crystal, m.p.  $147^\circ\text{C}$ .  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ , 400 MHz): 1,1 (6H, t,  $J=7.2$  Hz,  $\text{CH}_3$ ); 3,2 (4H, m,  $\text{CH}_2$ ); 7,3 (10H, m).  $^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ , 400 MHz) : 13,47 ( $\text{C}_8$ ); 42,99 ( $\text{C}_7$ ); 127,65 ( $=\text{C}_4$ ); 128,67 ( $=\text{C}_5$ ); 132,21 ( $=\text{C}_6$ ); 136,03 ( $=\text{C}_3$ ); 159,03 ( $\text{C}_1=\text{O}$ ); 170,96 ( $-\text{C}_2=\text{O}$ ). IR (KBr),  $\nu_{\text{maks}}$  ( $\text{cm}^{-1}$ ) : 3087 ( $\text{Csp}_2\text{-H}$ ), 2977 ( $\text{C-H}$ ), 1717 ( $\text{C=O}_{\text{ar}}$ ), 1676 ( $-\text{C=O}(-\text{N}-)$ ), 1580 ( $\text{C=C}$ ), 858 ( $\text{C-H}_{\text{ar}}$ ). MS (EI):  $m/z$  324 ( $\text{M}^+$ ), calculated Mass  $\text{C}_{19}\text{H}_{20}\text{N}_2\text{O}_3$ : 324.15.

The synthesis procedure used was capable to produce the designed compound. The main feature for the formation of the compound is the absence of hydrogen on both N of diethylurea and the presence of 2 benzoyl group. This is proved by  $^1\text{H-NMR}$  spectrum showing an additional H peak in the benzene region ( $\delta=7,3$  ppm). The  $\text{C=O}$  of benzoyl is proved by  $^{13}\text{C-NMR}$  chemical shift at 170 ppm, supported by the IR spectrum showing absorption band at  $1717\text{ cm}^{-1}$  and  $1676\text{ cm}^{-1}$ .

The *N,N'*-carbonylbis(*N*-ethylbenzamide) was produced by nucleophilic acylation [27] between benzoyl chlorides and *N,N'*-diethylurea. The structural change of *N,N'*-diethylurea to *N,N'*-carbonylbis(*N*-ethylbenzamide) was characterized by the conversion of  $-\text{NH}-$  moiety of diethylurea to  $-\text{N}-\text{C=O}(\text{ar})$ . As there are two symmetric  $-\text{NH}-$  moiety, the new compounds bear two symmetric benzoyl group, each attach on their N atom.

**Antitumor Activity:** The synthesized compound was subjected to *in vitro* antitumor activity test on human breast carcinoma cell line, compared with HU and 5-FU. The results were listed in Table 1. Table 1 show that *N,N'*-carbonylbis (*N*-ethylbenzamide) have  $IC_{50}$  lower than hydroxyurea and 5-FU. This indicated that the compound had antitumor activity against MCF-7 and T47D cells higher than HU. Its activity on MCF-7 also higher than 5-FU, an anticancer drug that is clinically used in the treatment of human breast carcinoma. It seems that benzoyl and ethyl group on the structure of urea may increase antitumor activity.

**Molecular Docking:** The capability of tested compound to interact with target molecule 3HEG was investigated by docking study, and the results were compared to HU. Human breast cancer were known to overexpress in MAP kinase, represented by 3HEG in this study. It was reported that HU could interacted with ERK, one type of this enzyme [29]. *N,N'*-carbonylbis(*N*-ethylbenzamide) display better score ( $\text{RS}=-74.55$ ) than HU ( $\text{RS}=-37.12$ ) which indicate high affinity of the new compound to 3HEG. Further visual inspection denoted that the carbonyl ( $\text{C=O}$ ) group and two N atom of *N,N'*-carbonylbis(*N*-ethylbenzamide) capable to form hydrogen bonds with amino acid residue Asp 168 in binding site of 3HEG (cavity-1). There is no hydrogen bond between HU and 3HEG. These interaction denoted the importance of  $\text{N-CO-N}$  moiety of urea for binding and the subsequent inhibitory capacity. Besides hydrogen bonds, there are van der Waals intermolecular forces (hydrophobic interactions) performed by compound with amino acid residues Asp 168, Tyr 35, Glu 71, Leu 167, and Phe 169. Such interactions supported the better score (RS) for the compound compare with the HU. Molecular docking of *N,N'*-carbonylbis(*N*-ethylbenzamide) to 3HEG is shown in Figure 2 and Figure 3. The comparative docking study of the *N,N'*-carbonylbis(*N*-ethylbenzamide) with anticancer drug hydroxyurea supports the molecular design of this new compound.


## CONCLUSION

of *N,N'*-carbonylbis(*N*-ethylbenzamide) has been designed and successfully synthesized from *N,N'*-

diethylurea and benzoyl chloride. The compound showed higher activity against human breast cancer cell lines (MCF-7 and T47D) than hydroxyurea. The compound was also more active against MCF-7 compared with 5-fluorouracil, an anticancer drug which has been clinically used in the treatment of human breast cancer. It is recommended to develop the compound as antitumor agents for human breast cancer.

#### Acknowledgement

We are thankful to The Indonesian Higher Education Directorate for financial support (PUPT/Prime Research Grant for University in 2014). We also thank to Institute for Tropical Disease Airlangga University and Faculty of Medicine Gajahmada University for their supports in this research.




**Figure 1:** Structure of *N,N'*-carbonylbis(*N*-ethylbenzamide)


**Table 1.** IC<sub>50</sub> growth inhibition activity of tested compounds against human breast cancer cell lines

Compound	IC <sub>50</sub> (mM)	
	MCF-7	T47D
<i>N,N'</i> -carbonylbis( <i>N</i> -ethylbenzamide)	0.64	0.79
Hydroxyurea	31.32	5.60
5-Fluorouracil	15.37	*

\* = not determined



**Figure 2 :** docked conformation of *N,N'*-carbonylbis(*N*-ethylbenzamide) molecule in secondary structure of p38α MAPK (2A), and in hydrophobic environment of enzyme's binding site, cavity-1 (2B)



**Figure 3 :** docked conformation showing hydrogen bonds between *N,N'*-carbonylbis(*N*-ethylbenzamide) molecule and amino acid residue Asp 168 (3A), and 2D-picture showing hydrophobic interaction performed by compound with amino acid residues Asp 168, Tyr 35, Glu 71, Leu 167, and Phe 169 (3B).

## REFERENCES

1. Stewart B, Wild C. Wild World Cancer Report, IARC Nonserial Publication, WHO Press: Lyon, 2014.
2. Goldman B. Multidrug resistance: can new drugs help chemotherapy score against cancer?. J. Natl. Cancer Inst. 2003; 95(4): 255–257.
3. Tibes R, Mesa RA. Blood consult: resistant and progressive essential thrombocythemia. Blood 2011; 118(2): 240–242.
4. Tartarone A et al. Mechanisms of resistance to EGFR tyrosine kinase inhibitors gefitinib/erlotinib and to ALK inhibitor crizotinib. Lung Cancer 2013; 81(3): 328–336.
5. Matsumura F. Studies on the action mechanism of benzoylurea insecticides to inhibit the process of chitin synthesis in insects: A review on the status of research activities in the past, the present and the future prospects, Pesticide Biochemistry and Physiology 2010; 97(2): 133–139.
6. Ramesh DK et al. Synthesis, Characterisation And Anticonvulsant Activity Of 1-(Aryl)-3- (Diphenylmethyl) Urea Derivatives. International Journal Of Research In Pharmacy And Chemistry (IJRPC) 2013; 3(4): 748–752.
7. Zheng Q-Z et al. Synthesis of some N-alkyl substituted urea derivatives as antibacterial and antifungal agents, European Journal of Medicinal Chemistry 2010; 45: 3207–3212.
8. Luzina EL, Popov AV. Anticancer activity of *N*-bis(trifluoromethyl)alkyl-*N'*-(polychlorophenyl) and *N'*-(1,2,4-triazolyl) ureas, European Journal of Medicinal Chemistry 2010; 45: 5507–5512.
9. Li Y et al. Discovery of novel urea derivatives as dual-target hypoglycemic agents that activate glucokinase and PPAR $\gamma$ . Eur J Med Chem. 2014; 9(76): 182–92.
10. Engen W et al. Synthesis of aryl-heteroaryl ureas (AHUs) based on 4-aminoquinoline and their evaluation against the insulin-like growth factor receptor (IGF-1R), Bioorganic & Medicinal Chemistry 2010; 18: 5995–6005.
11. Song EY et al. Synthesis of amide and urea derivatives of benzothiazole as Raf-1 inhibitor. European Journal of Medicinal Chemistry 2008; 43: 1519–1524.
12. Getlik M et al. Structure-based design, synthesis and biological evaluation of *N*-pyrazole, *N'*-thiazole urea inhibitors of MAP kinase p38, European Journal of Medicinal Chemistry 2012; 48: 1–15.
13. Wagner EF, Nebreda AR. Signal integration by JNK and p38 MAPK pathways in cancer development, Nat. Rev. Cancer 2009; 9: 537–549.
14. Harrison CN. Hydroxyurea compared with anagrelide in high-risk essential thrombocythemia. N. Engl. J. Med. 2005; 353(1): 33–45.
15. Escibano L et al. Prognosis in adult indolent systemic mastocytosis: A long-term study of the Spanish Network on Mastocytosis in a series of 145 patients. Journal of Allergy and Clinical Immunology. 2009; 124 (3): 514–521.
16. Barosi G et al. A unified definition of clinical resistance/intolerance to hydroxyurea in essential thrombocythemia: results of a consensus process by an international working group. Bioorganic & Medicinal Chemistry Letters 2009; 19: 755–758.
17. Rajwade RP et al. Hydroxyamic Acid Analogous Against Breast Cancer Cells: 2D QSAR and 3D QSAR Studies. Molecular Informatics (accessed December 27, 2014)
18. Avendano C, Menendes JC. Medicinal Chemistry of Anticancer Drugs. Elsevier: Amsterdam, 2008. pp. 13-18.
19. Nambodiri HV et al. Analysis of imatinib and sorafenib binding to p38alpha compared with c-Abl and b-Raf provides structural insights for understanding the selectivity of inhibitors targeting the DFG-out form of protein kinases. Journal Biochemistry 2010; 49: 3611–3618.
20. Saban N, Bujak M. Hydroxyurea Hydroxamic Acids and Derivatives as Antitumor Drugs. Cancer Chemother Pharmacol 2009; 64: 213–221.
21. Diyah NW et al. Molecular modeling and Quantitative Structure-Activity Relationship of Benzoylurea as Antitumor. Pharmaceutical Chemistry Bulletin 2013, 2(2): 20–27 [in Bahasa Indonesia].
22. Diyah NW. In: Synthesis, Molecular Docking, and Antitumor Activity of *N,N'*-Dibenzoyl-*N,N'*-Dimethylurea against Human Breast Cancer Cell Line (MCF-7), Proceedings of the First International Conference on Pharmacy and Pharmaceutical Sciences,

- Surabaya, Indonesia, November 14–15, 2014; Wouter LJ Hinrichs, Kozo Takayama, Srinivas Mutalik, Nuttakorn Baisaeng, Widji Soeratri, Eds; Faculty of Pharmacy Airlangga University, Indonesia, 2014; pp. 203–205
23. Li JN *et al.* Inhibition of Microtubule Polymerization by 3-Bromopropionylamino-benzoylurea (JIMB01), a New Cancericidal Tubulin Ligand. *Biochemical Pharmacology* 2003; 65: 1691–1699.
  24. Fortin S *et al.* *N*-Phenyl-*N'*-(2-chloroethyl)ureas (CEU) as potential antineoplastic agents. Part 2: Role of  $\omega$ -hydroxyl group in the covalent binding to  $\beta$ -tubulin. *Bioorganic & Medicinal Chemistry* 2007; 15: 6703–6712.
  25. El-Sawy E *et al.* Synthesis antimicrobial and anticancer activities of some new *N*-ethyl, *N*-benzyl and *N*-benzoyl-3-indolyl heterocycles, *Acta Pharm* 2012, 62: 157–179.
  26. Lu C-S *et al.* Synthesis and in vitro antitumor activities of novel benzylurea analogues of sorafenib, *Acta Pharmaceutica Sinica* 2013, 48(5): 709–717.
  27. McMurry JM. *Organic Chemistry with Biological Application*, 2<sup>nd</sup> ed.; Brooks/Cole: Belmont, 2011.
  28. RCSB Protein Data Bank. <http://www.rcsb.org/pdb> (Accessed February 23, **2015**)
  29. Wei F *et al.* ERK1 and ERK2 kinases activate hydroxyurea-induced S-phase checkpoint in MCF7 cells by mediating ATR activation. *Cellular Signalling* 2011; 23: 259–268.

# World Journal of Pharmaceutical Sciences



[Register](#) [Login](#)



ISSN (Print): 2321-3310

ISSN (Online): 2321-3086

US CODEN : WJPSBT

\* Impact Factor: 1.877 (2013)



## Editorial Board

### Editor-in-Chief

**Dr. Kiran K. Akula**

Department of Biology & Biochemistry

University of Houston

**USA**

## Executive Editors

**Dr. Dennis K. Miller,**  
Department of Psychological  
Sciences,  
University of Missouri,  
**USA**

**Dr. Khaled Nabih Zaki Rashed,**  
Pharmacognosy Department,  
Pharmaceutical and Drug Industries Research  
Division,  
**Egypt**

## Associate Editors

**Prof. Jinyong Peng,**  
College of Pharmacy,  
Dalian Medical University,  
**China**

**Prof. Chika Jhon Mbah,**  
Faculty of Pharmaceutical Sciences,  
University of Nigeria,  
**Nigeria**

**Prof. Shaker A. Mousa,**  
Vice Provost for Research,  
Albany College of Pharmacy and Health  
Sciences,  
**USA**

**Dr. Moawia M. A. M. Altabakha,**  
College of Pharmacy,  
Al Ain University of Science and  
Technology,  
**UAE**

## Editorial Members

**Dr. Eyad S. M. Abu-Nameh, Jordan**

**Dr. Fadi M. Alkhateeb, USA**

**Dr. Babak Nahri-Niknafs, Iran**

**Dr. Md. Mustafizur Rahman, Bangladesh**

**Dr. S. N. Manjula, India**

**Dr. Rohini Karunakaran, Malaysia**

**Dr. Wael abu Dayyih, Jordan**

**Dr. Ravi S. Shukla, USA**

**Dr. Rashad Mohammed Musleh Alnamer, Yemen**

**Dr. Mosad A. Ghareeb, Egypt**

**Dr. Muhammad Hanif, Pakistan**

**Dr. Venkata Srikanth Meka, Malaysia**

**Dr. Priyanka M. Jadhav, USA**

**Dr. Sumi Mukhopadhyay, India**

**Dr. Mahtab Z. Siddiqui, India**

**Dr. Laleh Khodaie, Iran**

**Prof. Mamdouh Moawad Ali, Egypt**

**Dr. G. Narasimha, India**

**Dr. Diana Malaeb, Lebanon**

**Dr. Ashaben Patel, USA**

**Dr. Mario Ariatti, South Africa**

**Prof. Eric de Souza Gil, Brazil**

**Dr. Sadia Sultan, Malaysia**

**Dr. Hanan Abd El-Monem Ahmed Merey, Egypt**

**Dr. Milind Sadashiv Alai, Taiwan**

**Dr. Shazia Jamshed, Malaysia**

**Dr. Victoria Cecilia Cristea, Romania**

**Prof. Dhrubo Jyoti Sen, India**

**Dr. Binaya Sapkota, Nepal**

**Dr. Jagdish L. Kakadiya, India**

**Dr. Mohsen Mohamed Selim Asker, Egypt**

**Harsimran Singh Bhalla, India**

**Dr. R. Kavitha, India**

**Dr. Neslihan Üstündag Okur, Turkey**

**Dr. Basavaraj C. Metri, India**

**Prof. Dr. Mohamed Salama Mohamed, Malaysia**

**Dr. Nisha Mary Joseph, Ethiopia**

**Dr. Sujit Kumar Kar, India**

**Dr. Gamal Osaman Elhassan, Saudi Arabia**

**Dr. Syed Ahmed Iizhar, Saudi Arabia**

**Md. Shafayat Hossain, China**

**Dr. Mohd Masih Uzzaman Khan, Saudi Arabia**

**Dr. Nagib A. Elmarzugi, Malaysia**

**Dr. Yasser M.S.A. Al-Kahraman, Pakistan**

**Ruth Elenora Kristanty, Indonesia**

**Dr. Theurkar Sagar Vasant, India**

**Dr. Pramod Singh Khatri, India**

**Dr. Vahid Rakhshan, Iran**

**Dr. Shashank Kumar, India**

**Prof. Rajni Bala, India**

**Dr. Javad S. Rad, Iran**

**Dr. Sandeep Kumar Gupta, India**

**Dr. Narendra Kumar, India**

**Dr. DSVGK Kaladhar, India**

**Dr. Ofentse Mazimba, Botswana**

**Dr. Udit Tiwari, India**

**Ari Yuniarto, Indonesia**

**Prashant Kale, India**

**Prof. J.F. Eniojukan, Nigeria**

**Dr. Jigna G. Tank, India**

**Dr. Behzad Foroutan, Iran**

**Dr. Sarika Singh, India**

**Dr. Atta Abbas, UK**

## Archives

[Volume-1 \(2013\)](#)  [Volume-2 \(2014\)](#)  [Volume-3 \(2015\)](#) 

## Latest Updates

- 
- 
- 

## Manuscript Status

**Enter your Manuscript Number to know status of your Manuscript**



## Abstracted and Indexed In







A division of the American Chemical Society